



| | | | | | | | | | |
|--------|------------|---------|---------------|-----------|-----------|----------|------|-------|-------|
| PubMed | Nucleotide | Protein | Genome | Structure | PopSet | Taxonomy | OMIM | Books | |
| Search | PubMed | ▼ | for | | | | | Go | Clear |
| | | Limits | Preview/Index | History | Clipboard | Details | | | |

| | | | | | | | | |
|---------|----------|---|------|---|------|------|------------------|-------|
| Display | Abstract | ▼ | Sort | ▼ | Save | Text | Add to Clipboard | Order |
|---------|----------|---|------|---|------|------|------------------|-------|

1: J Clin Microbiol 1990 Mar;28(3):495-503

Related Articles, Boo

Rapid and simple method for purification of nucleic acids.

Boom R, Sol CJ, Salimans MM, Jansen CL, Wertheim-van Dillen PM, van de Noordaa J.

Department of Virology, Academic Medical Center, Amsterdam, The Netherlands.

We have developed a simple, rapid, and reliable protocol for the small-scale purification of DNA and RNA from, e.g., human serum and urine. The method is based on the lysing and nuclease-inactivating properties of the chaotropic agent guanidinium thiocyanate together with the nucleic acid-binding properties of silica particles or diatoms in the presence of this agent. By using size-fractionated silica particles, nucleic acids (covalently closed circular, relaxed circular, and linear double-stranded DNA; single-stranded DNA; and rRNA) could be purified from 1: different specimens in less than 1 h and were recovered in the initial reaction vesse Purified DNA (although significantly sheared) was a good substrate for restriction endonucleases and DNA ligase and was recovered with high yields (usually over 50%) from the picogram to the microgram level. Copurified rRNA was recovered almost undegraded. Substituting size-fractionated silica particles for diatoms (the fossilized cell walls of unicellular algae) allowed for the purification of microgram amounts of genomic DNA, plasmid DNA, and rRNA from cell-rich sources, as exemplified for pathogenic gram-negative bacteria. In this paper, we show representative experiments illustrating some characteristics of the procedure which may have wide application in clinical microbiology.

PMID: 1691208 [PubMed - indexed for MEDLINE]

| | | | | | | | | |
|---------|----------|---|------|---|------|------|------------------|-------|
| Display | Abstract | ▼ | Sort | ▼ | Save | Text | Add to Clipboard | Order |
|---------|----------|---|------|---|------|------|------------------|-------|

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer